Initiated: 5/12/04

Reviewed/revised: 8/1/13

Patient with any of the following:

Intubation/assisted ventilation


Revision: 3

MILWAUKEE COUNTY EMS STANDARD OF CARE TRAUMA TEAM ACTIVATION -

Approved by: M. Riccardo Colella, DO, MPH, FACEP

Page 1 of 1

PEDIATRIC PATIENTS



CRITERIA:

Breathing: Respiratory arrest

Respiratory distress (ineffective respiratory effort, stridor or grunting)

Respiratory Rate:

0-5 months: RR less than 20 6 months -12 years: RR less than 16 13 years or older: RR less than 12

Circulation:

0 - 5 months: Systolic BP less than 60 or Pulse less than 90

6 months - 5 years: Systolic BP less than 70 or Pulse less than 70

6 years or older: Systolic BP less than 90 or Pulse less than 60

Clinical signs of shock:

Pale, cold, clammy, tachycardia with weak pulses, capillary refill greater then 3 seconds in a warm environment

Patients receiving blood Patients receiving more than 40 mL/kg fluids

Neurological: GCS less than 13

Anatomic Diagnosis:

Penetrating injury to head, neck, torso, groin Flail chest

Open chest wound

Level I Trauma

Open or depressed skull fracture Suspected spine or spinal cord injury Bilateral femur fractures

Pelvic fracture

Suspected airway injury from facial burns, chemical or smoke inhalation

Complete/partial amputation or degloving above wrist or above ankle

Deep penetrating injuries above knee or above elbow

Significant blunt maxillofacial trauma

Level I I Trauma

mechanisms of injury, who does not meet the Level I or Level II criteria:

Crash speed more than 20 mph Passenger ejected from vehicle Vehicle roll-over Death or severe injury of same-car

occupant Pedestrian hit by vehicle and thrown, dragged or run over

Pedestrian struck at 20 mph or more Passenger compartment intrusion more than 12"

Prolonged extrication - over 20 minutes

Extremity injury with concern for ischemia

Trauma Consult